

LETTER TO THE EDITOR

Open Access

A hypothesis on chemical mechanism of the effect of hydrogen

Penghui Shi^{1,3*†}, Wancang Sun^{1†} and Pengzhong Shi²

Abstract

Many studies have shown that hydrogen can play important roles on the antioxidant, anti-inflammatory and other protective effects. Ohsawa et al have proved that hydrogen can electively and directly scavenge hydroxyl radical. But this mechanism cannot explain more new experimental results. In this article, the hypothesis, which is inspired by H₂ could bind to the metal as a ligand, come up to explain its extensive biology effect: Hydrogen could regulate particular metalloproteins by bonding (M-H₂ interaction) it. And then it could affect the metabolization of ROS and signal transduction. Metalloproteins may be ones of the target molecules of H₂ action. Metal ions may be appropriate role sites for H₂ molecules. The hypothesis pointed out a new direction to clarify its mechanisms.

Keywords: Hypothesis, Hydrogen, Mechanism, Ligand

Background

Basal cellular metabolism continuously produces reactive oxygen species (ROS). O₂ may generate successively superoxide (O₂⁻), hydrogen peroxide (H₂O₂) and hydroxyl radical (OH[·]). ROS are able to oxidize biological macromolecules such as DNA, protein and lipids [1,2]. Some enzymic systems detoxify ROS, however, catalase dismutates H₂O₂, and SOD eliminates O₂⁻ (but generates H₂O₂). Excess oxygen can react with H₂O₂ to produce hydroxyl radicals by the Fenton reaction [3] (Figure 1). Ohsawa et al provide evidence that hydrogen could reach subcellular compartments such as the nucleus and mitochondria, biochemical experiments using fluorescent probes and electron paramagnetic resonance spectroscopy spin traps indicated that hydrogen gas may selectively scavenge the hydroxyl radical [4]; They were the first to show the ability of H₂ to suppress oxidation in vivo. So far, many researches have proved the central role of hydrogen on the antioxidant, anti-inflammatory and other protective effects.

Much evidence has shown that hydrogen exert beneficial effects in animal models of a number of diseases mainly associated with oxidative stress; However, the findings don't cover plants. Moreover, the mechanism of the effect of hydrogen remains unclear, the most acceptable mechanism is that the hydrogen can electively and directly scavenge hydroxyl radical while preserving other reactive oxygen and nitrogen species important in signaling [4]. There is some question, however, the published rate constant for the reaction of ·OH with H₂ to form H₂O and H[·] is drastically slower than most radical-radical reactions [6]. Furthermore, it can't explain some new experimental results. For example, Tomohiro Itoh proved that hydrogen exerts its beneficial effect by modulating some signaling pathways. Experimenters found that oral intake of hydrogen-rich water abolishes an immediate-type allergic reaction in mice. The results indicated that hydrogen attenuates phosphorylation of the FcεRI-associated Lyn and its downstream signal transduction, which subsequently inhibits the NADPH oxidase activity and reduces the generation of hydrogen peroxide. they also found that inhibition of NADPH oxidase attenuates phosphorylation of Lyn in mast cells, indicating the presence of a feed-forward loop that potentiates the allergic responses. Hydrogen accordingly inhibits all tested signaling molecule(s) in the loop. The results imply that effects of hydrogen in some diseases are possibly mediated by modulation of yet unidentified signaling pathways [7].

* Correspondence: shiph150@sina.cn

†Equal contributors

¹Gansu Key Lab of Crop Improvement & Germplasm Enhancement, Gansu Provincial Key Laboratory of Aridland Crop Sciences, Gansu Agricultural University, Lanzhou 730070, China

³Gansu Agricultural University, Anning District, Lanzhou City, Gansu Province, China

Full list of author information is available at the end of the article





Figure 1 ROS generation and detoxification. Various chemical reactions, with or without enzymic catalysis, generate ROS. The dioxygen molecule undergoes successive reductions which yield the superoxide radical anion ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2) and the hydroxyl radical ($OH\cdot$). Antioxidant systems act as ROS scavengers to maintain the intracellular redox status. Quinone reductase (QR) detoxifies quinone compounds, metallothionein (MT) traps (heavy) metal cations, and vitamins C and E trap free radicals. SOD and catalase respectively dismutate superoxide (into oxygen and hydrogen peroxide) and hydrogen peroxide (into oxygen and water). Glutathione peroxidase (GPx) acts like catalase on various peroxide compounds, including H_2O_2 [5].

Hypothesis and discussion

Although the beneficial effect of hydrogen is generally accepted, the mechanism is not still clear. There can be little doubt that biology function of H_2 depends on the physical and chemical interaction of other molecules with it, so what would it be like?

In organometallic and inorganic chemistry, for some metal complexes, the "arrested" addition product can be isolated—the dihydrogen complex is obtained as a stable species that can be put in a bottle: [8] (Scheme 1).

Traditional ligands use lone pairs of electrons in their bonding, but in dihydrogen complexes, the bonding to the metal comes from donation of electron density from the nonpolar H-H σ bond to d orbitals on the metal (Figure 2). We hold that hydrogenase is a typical case in point. Hydrogenases catalyse the reversible oxidation of molecular hydrogen (H_2). The active site domain of the Fe hydrogenases contains an unusual Fe-S centre termed the H-cluster. H-cluster consists of the $[Fe_4S_4]$ subcluster bridged via the Cys thiolate to the $[Fe_2]$ (binuclear iron) subcluster (Figure 3) [9-12]. Iron sulfur clusters are found at the active sites of numerous enzymes where they commonly facilitate electron transfer and substrate transformations (Table 1) [13]. It is inferred that M- H_2 interaction also exists in these metalloproteins. By extension, it also should exist in non-cluster metalloproteins. All this suggests that metal ions may be the site where H_2 interacted with metalloproteins.



Scheme 1 Formation of a dihydrogen complex.

It is proved from experiments that molecules containing a metallic cation may promote O_2^- formation because they have the ability to store and easily give an electron to molecular dioxygen [16]. Free radicals arise through the autoxidation catalyzed by metalloproteins, this mainly occurs within the mitochondria. Some experimental results proved that hydrogen can permeate




Figure 2 Schematic bonding model between molecular hydrogen and a metal [14].




Figure 3 Stereo view of the Cpl H-cluster and coordinating Cys ligands located at the boundary of the two lobes of the active site domain [15].

into mitochondria and prevent superoxide formation [17]. It indicates that H_2 can affect the metabolization of ROS by way of superoxide formation. And it must have a big impact on the content of H_2O_2 . Besides, free radical production can also happen in other cellular compartments, such as NADPH oxidase. ROS production can interfere with signal transduction pathways [18-20]. ROS, in particular H_2O_2 , are indeed second messengers for many physiological stimuli, some stimuli have been proven to induce mitochondrial H_2O_2 release [21].

On the basis of the mentioned analysis, we evaluate that metal ions could be appropriate role sites for H_2 molecules. We propose that hydrogen can permeate into mitochondria and concentrate at mitochondrial membrane to regulate activity of metalloproteins (complexes I, II, III) by M- H_2 interaction. The same may be true of NADPH oxidase. This could reduce the production of superoxide and prevent synthesis of the hydroxyl from the source. Hydrogen, on the one hand, maybe regulate the content of H_2O_2 by affecting the metabolization of ROS. And then it disturbs signaling pathway. On the other hand, hydrogen maybe directly influence signal transduction by regulating particular metalloproteins of signaling pathways. It may be the mechanism of its extensive biology effect.

Conclusions

We propose that metalloproteins may be ones of the target molecules of H_2 action. Metal ions may be appropriate role sites for H_2 molecules. And in this way can we explain

Table 1 Iron-sulphur proteins in cells

Function	Protein class
Catalysis	Bacterial nitrate reductase, Formate dehydrogenase, Fumarate reductase, Glutamine PRPP amidotransferase, Hydrogenase, Methane monooxygenase, NADH: ubiquinone reductase, Phthalate dioxygenase reductase, Succinate dehydrogenase, Sulphite reductase, Xanthine dehydrogenase, Aconitase (TCA cycle)
Electron transfer	Ferredoxins, Rieske proteins, Rubredoxins, NADH Dehydrogenase, Succinate-CoQ Reductase, CoQ-cyt c Reductase (respiratory chain complexes)

its extensive biology effect. Although some details remain murky, the hypothesis pointed out a new direction for the continuation. It is a good inspiration to clarify the mechanism of the effect of hydrogen. We predict that hydrogen can affect many metalloproteins activities. Therefore, more studies will be necessary to test the hypothesis.

Acknowledgment

We wish to sincerely thank Xuejun Sun (Second Military Medical University, Shanghai, China) for advice about the manuscript.

Funding

There is no source of support in the form of grants.

Author details

¹Gansu Key Lab of Crop Improvement & Germplasm Enhancement, Gansu Provincial Key Laboratory of Aridland Crop Sciences, Gansu Agricultural University, Lanzhou 730070, China. ²Inner Mongolia University of Science and Technology, School of Material and Metallurgy, Baotou 014010, China. ³Gansu Agricultural University, Anning District, Lanzhou City, Gansu Province, China.

Received: 23 May 2012 Accepted: 23 May 2012

Published: 21 June 2012

References

- Vaughan M: Oxidative modification of macromolecules minireview series. *J Biol Chem* 1997, **272**:18513.
- Hagen TM, Ingersoll RT, Lykkesfeldt J, Liu J, Wehr CM, Vinarsky V, Bartholomew JC, Ames AB: (R)-alpha-lipoic acid-supplemented old rats have improved mitochondrial function, decreased oxidative damage, and increased metabolic rate. *FASEB J* 1999, **13**:411-418.
- Woo ES, Lazo JS: Nucleocytoplasmic functionality of metallothionein. *Cancer Res* 1997, **57**:4236-4241.
- Ohsawa I, Ishikawa M, Takahashi K, et al: Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals. *Nat Med* 2007, **13**:688-694.
- Morel Y, Barouki R: Repression of gene expression by oxidative stress. *Biochem J* 1999, **342**:481-496.
- Wood KC, Gladwin MT: The hydrogen highway to reperfusion therapy. *Nat Med* 2007, **13**:673-674.
- Itoh T, Fujita Y, Ito M, et al: Molecular hydrogen suppresses FcεRI-mediated signal transduction and prevents degranulation of mast cells. *Biochem Biophys Res Commun* 2009, **389**:651-656.
- Kubas GJ, Ryan RR, Swanson BI, Vergamini PJ, Wasserman HJ: *J Am Chem Soc* 1984, **106**:451-452.
- Darensbourg MY, Lyon EJ, Smee JJ: The bio-organometallic chemistry of active site iron in hydrogenases. *Coord Chem Rev* 2000, **219/221**:533-561.
- Marr AC, Spencer DJE, Schurder M: Structural mimics for the active site of [NiFe] hydrogenase. *Coord Chem Rev* 2001, **219/221**:1055-1074.
- Nicolet Y, de Lacey AL, Vernde X, et al: Crystallographic and FTIR Spectroscopic Evidence of Changes in Fe Coordination Upon Reduction

of the Active Site of the Fe-Only Hydrogenase from *Desulfovibrio desulfuricans*. *J Am Chem Soc* 2001, **123**:1596–1601.

12. Lawrence JD, Li HX, Rauchfuss TB, *et al*: *Angew Chem Int Ed* 2001, **40**:1768–1771.
13. Cowan JA: *Inorganic Biochemistry: An Introduction*. New York: VCH Publishers; 1993.
14. Martin HG: *PrechtI*. WiKu: Publisher; 2007.
15. He Chengjiang WangMei, Li Minna, Sun Licheng: **Advance in chemical mimic of fe-only hydrogenase**. *Progress in chemistry* 2004, **16**(2):250–255.
16. Kagan VE, Tyurina YY: **Recycling and redox cycling of phenolic antioxidants**. *Ann N Y Acad Sci* 1998, **854**:425–434.
17. Sato Y, Kajiyama S, Amano A, *et al*: **Hydrogen-rich pure water prevents superoxide formation in brain slices of vitamin C-depleted SMP30/GNL knockout mice**. *Biochem Biophys Res Commun* 2008, **375**:346–350.
18. Finkel T: **Oxygen radicals and signaling**. *Curr Opin Cell Biol* 1998, **10**:248–253.
19. Lander HM: **An essential role for free radicals and derived species in signal transduction**. *FASEB J* 1997, **11**:118–124.
20. Suzuki YJ, Forman HJ, Sevanian A: **Oxidants as stimulators of signal transduction**. *Free Radical Biol. Med* 1997, **22**:269–285.
21. Quillet-Mary A, Jaffrezou JP, Mansat V, Bordier C, Naval J, Laurent G: **Implication of mitochondrial hydrogen peroxide generation in ceramide-induced apoptosis**. *J Biol Chem* 1997, **272**:21388–21395.

doi:10.1186/2045-9912-2-17

Cite this article as: Shi *et al*: A hypothesis on chemical mechanism of the effect of hydrogen. *Medical Gas Research* 2012 **2**:17.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

